

UNIVERSITY OF NAPLES FEDERICO II

DOCTORATE

MOLECULAR MEDICINE AND MEDICAL BIOTECHNOLOGY

XXX CICLO



**Integration of cAMP signaling and the ubiquitin system in the control of
primary cilium**

Tutor

Prof. Antonio Feliciello

Candidate

Simona Sauchella

COORDINATOR

Prof. Vittorio Enrico

Avvedimento

Academic Year 2016/2017

INDEX

ABSTRACT	1
1 INTRODUCTION	2
1.1 <i>The cAMP-dependent signal transduction pathway</i>	2
1.2 <i>Protein Kinase A (PKA)</i>	5
1.3 <i>AKAP proteins</i>	8
1.4 <i>The ubiquitin proteasome system and E3 ubiquitin ligase CHIP</i>	10
1.5 <i>Ciliogenesis</i>	13
1.6 <i>Correlation between cAMP signaling, the UPS system and primary cilium</i>	16
1.7 <i>NimA related kinase 10 (Nek10)</i>	18
2 AIM OF THESIS	21
3 MATERIALS AND METHODS	23
4 RESULTS	27
4.1 <i>NEK10, RIIβ and PCM1 form a macromolecular complex</i>	27
4.2 <i>Endogenous PCM1, NEK10 and RIIβ colocalize in Human Embryonic Kidney 293 cell</i> .	29
4.3 <i>NEK10 is required for ciliogenesis</i>	30
4.4 <i>PKA regulates the stability of primary cilium</i>	33
4.5 <i>PKA phosphorylation primes NEK10 for proteolysis via UPS</i>	35
4.6 <i>CHIP is the NEK10 E3 ubiquitin ligase</i>	40
4.7 <i>Dysregulation of CHIP affects cilia in SCAR16 disease</i>	44
5 DISCUSSION AND CONCLUSION	46
6 APPENDICES	51
7 ACKNOWLEDGEMENTS	52
8 REFERENCES	53
9 LIST OF PUBLICATIONS	57

ABSTRACT

The primary cilium is an antenna-like sensory organelle able to receive extracellular signals and it is localized on the surface of most human cells.

In my thesis, I investigated the connection between G-protein coupled receptor (GPCR) signaling and the ubiquitin proteasome system (UPS) pathway in the control of cilium stability. I identified, at pericentriolar region, a trimeric complex composed by PCM1, NEK10 and PKA. I demonstrated that NEK10 has a crucial role for ciliogenesis. Phosphorylation by PKA primes NEK10 to proteasomal degradation. Disappearance of NEK10 promotes cilia resorption. I identified CHIP as the E3 ubiquitin ligase responsible of NEK10 ubiquitination and I demonstrated that CHIP mediates the effects of cAMP on primary cilium stability.

Dearrangement of this control mechanism was observed in proliferative and genetic disorders. Collectively, the findings unveil a pericentriolar kinase signalosome that efficiently links the cAMP cascade with the ubiquitin-proteasome system, controlling essential aspects of ciliogenesis.

1 INTRODUCTION

1.1 The cAMP-dependent signal transduction pathway

The biological organisms are able to modify a variety of cellular processes to adapt themselves to multiple conditions. Subsequently to a change, the organism communicates to specific target cells through the extracellular messengers such as hormones, neurotransmitters and growth factors.

The binding between a ligand and its specific receptor on the cell surface can lead to a different biophysical response through activation of specific second messengers.

The second messengers trigger a biological response that may consist in the regulation of different cellular processes, as gene transcription, protein translation, hormone production, cellular differentiation and cellular division.

The cyclic AMP (cAMP) is the most famous second messenger because it is involved in a wide array of biological processes^{1,2}.

When an extracellular ligand binds a G-protein coupled receptors (GPCR), it starts the cAMP signaling cascade.


The G protein-coupled family receptors (GPCRs) are a large family of trans-membrane proteins that transduce extracellular signals into the cell³. The binding of extracellular ligand to its GPCR activates the adenylyl cyclase (AC), an enzyme that converts the ATP in cAMP⁴. The activity of ACs is stimulated by the interaction with the stimulatory α subunit of the G-protein ($G_{\alpha s}$). In basal conditions, $G_{\alpha s}$ forms an heterotrimeric complex with β and γ subunits. Subsequently to the binding of the extracellular messenger, the GPCRs causes the dissociation of heterotrimeric G-proteins, with consequent activation of ACs by the $G_{\alpha s}$ subunit⁵.

In addition to AC, the levels of cAMP are regulated by the cyclic nucleotide phosphodiesterases (PDEs) and phosphatases (PPs).

Phosphodiesterases (PDEs) are involved in the regulation of the intracellular level of cAMP. They are part of a large superfamily of enzymes that hydrolyze the 3' - 5' phosphodiester bond in the second messenger cAMP with the formation of 5'-AMP⁶.

By reducing the levels of cAMP, PDEs regulate the duration and amplitude of the cyclic nucleotide signaling⁷. The subcellular localization of the enzymes is controlled by the N-terminal regulatory region⁸. The distribution of PDEs in the cells generates intracellular micro domains that locally enhance the sensitivity and specificity of the intracellular response to the cAMP⁹.

There are three distinct classes of direct effectors of the cAMP : cAMP-dependent protein kinase (PKA), RAP exchange proteins (EPACs), and cAMP gated ion channels (cNGC) (**Fig.1**). Among these effectors, the more studied is Protein kinase A (PKA).



Nature Reviews | Molecular Cell Biology

Figure 1. Schematic diagram of cAMP synthesis and downstream effectors activation. When an extracellular ligand binds to and activates a seven-transmembrane G-protein-coupled receptor, the signal is passed through the heterotrimeric G protein to adenylyl cyclase. The activated adenylyl cyclase converts ATP into the second messenger cAMP. Principal effectors of cAMP are PKA, PDE and EPAC. The cyclic nucleotide phosphodiesterases (PDE) degrades the phosphodiester bond in the second messenger molecules cAMP and cGMP. RAP exchange proteins (EPACs) are cAMP-dependent guanine-nucleotide-exchange factors for the small GTPases, and are known to be important mediators of cAMP signaling. Finally there are cAMP gated ion channels (cNGC) that function in response to the binding of cyclic nucleotides as cGMP and cAMP.

1.2 Protein Kinase A (PKA)

Protein kinase A (PKA) is a serine/threonine kinase and it is one of the major effector of the cAMP. Indeed, every time an extracellular stimulus leads to an intracellular response cAMP-mediated, PKA is activated¹⁰.

PKA is a tetrameric protein composed by two catalytic (C) and two regulatory(R) subunits. The whole complex makes the holoenzyme inactive.

In response to the binding of a ligand to GPCRs, there is a quick increase of the intracellular concentration of cAMP generated by the ACs proteins². So cAMP binds to R subunits, this binding causes their dissociation from the catalytic subunits (**fig. 2**). In this way, the catalytic subunits are able to phosphorylate many different downstream cellular substrates that include ion channels and a lot of transcription factors².

The biochemical and functional features of PKA holoenzyme are largely determined by the structure, properties and relative abundance of the R subunits¹¹. The conserved catalytic core in the C-subunit is encoded from three different genes, C α , C β e C γ ¹². The catalytic subunit is a 350-amino acid protein and the kinase core is localized into 40-300 residues. The smaller N-terminal lobe is composed by β -sheets and is responsible for nucleotide binding¹³, whereas the larger C-terminal lobe is composed by α -helices and it is responsible of substrates binding and catalysis¹¹.

R-subunits are encoded by four different genes (R1 α , R1 β , R2 α e R2 β) that confer the different biochemical and biological characteristics to the PKA isoforms¹². The R-subunit polypeptide contains an NH₂-terminal dimerization domain, an autophosphorylation site (that is the principal contact site for the C subunit) and two cAMP binding sites. Another functional site present on the N-terminus of the R-subunits is the dimerization/docking (D/D) domain that provides a docking site

for the A Kinase Anchoring Proteins (AKAPs) ¹⁴. In addition, the R subunits are able to form both homo- and heterodimers generating a large number of combinations, which further contribute to diversity and presumably specificity in the cAMP signal pathway¹⁵.

The PKAs that contain either RI or RII are identified as PKA type I or type II and they have different sensitivities to cAMP. They also differ for localization and expression. PKA type I, in fact, is largely cytoplasmic, whereas PKA type II is confined to subcellular structures and compartments. Furthermore RI α and RII α are ubiquitously expressed, RI β has been mainly abundant in neuronal tissues while RII β has the highest expression in neuronal, adipose, testes and heart tissues^{12, 16}. Studies demonstrated that ablation of the gene encoding the RI β leads to deficits in hippocampal long-term depression and depotentiation^{17, 18} but with a compensatory increase in total PKA activity, suggesting a unique role for RI β in synaptic plasticity¹⁹. A targeted disruption of the RII α gene yields viable mice with no physiological abnormalities, implying that PKAI and/or PKAII β compensates for the RII α defect^{20, 21}. The mutant mice with disruption of the mouse RII β gene are lean and have elevated metabolic rates caused by increases in both basal PKA activity and the basal rate of lipolysis^{22, 23}. RII β KO mice also display defects in neuronal gene expression, learning and behavior^{24, 25}. The activity of PKA is regulated by specific protein phosphatases. It has been demonstrated that phosphatases belonging to the PP1 and PP2A families are responsible for dephosphorylation of PKA substrates. In turn, PKA can control phosphatase activity by phosphorylation of specific PP1 inhibitors, such as I-1 and DARPP32²⁶.

PKA signaling is compartmentalized thanks to AKAPs protein. They are a group of several scaffold proteins that anchor the R subunits to tissues and different cellular compartments.¹⁵




Figure 2. PKA molecular structure and activation mechanism.

When two molecules of cAMP bind the regulatory subunits of PKA, the holoenzyme is disassociates causing the activation of catalytic subunits.

1.3 AKAP proteins


PKA is targeted to specific cellular organelles or subcellular locations through interaction with a family of distinct but functionally homologous proteins called AKAPs (A-Kinase-Anchor-Proteins)²⁷⁻²⁹.

AKAPs contain a PKA-binding motif, approximately 14 amino acids, that are able to bind the R subunit^{7, 30}. This region forms an amphipathic helix in which hydrophobic residues are located in the interior face while charged residues align on the exterior surface. This helical wheel binds with high affinity the N-terminal docking/dimerization (D/D) domain of the PKA-R dimer directing it in proximity of its substrates³¹ (**Fig. 3**). In the past have been identified several AKAPs that bind both RI and RII subunits³².

Through their targeting domain, AKAPs protein have been found in various cellular organelles such as centrosomes, dendrites, endoplasmic reticulum, mitochondria, nuclear membrane, plasma membrane and vesicles. The presence of PKA nearby of its substrates enhances the PKA-dependent phosphorylation of a large number of cellular substrates. In fact the cells that express high levels of PKA are more responsive to signals caused by the intracellular increase of second messenger cAMP^{33, 34}.

Although AKAPs have been defined on the basis of their interaction with PKA, several of these molecules are able to bind other enzymes such as receptors, effectors, protein phosphatases and kinases. In fact in the cells, AKAPs form macro molecular complexes, named transduceosome, where distinct signaling pathways converge and are attenuated or amplified, improving the specificity and efficiency of biological responses³⁴⁻³⁶.

S-AKAP84	E	I	K	R	A	A	F	Q	I	I	S	Q	V	I	S	E	A	T
Ht-31	L	I	E	E	A	A	S	R	I	V	D	A	V	I	E	Q	E	K
AKAP350	V	E	E	K	V	A	A	A	L	V	S	Q	I	Q	L	E	A	V
AKAP150	L	L	I	E	T	A	S	S	L	V	K	N	A	I	E	L	S	V
AKAP95	T	P	E	E	V	A	A	E	V	L	A	E	V	I	T	A	A	V
YOT1AO	R	L	E	E	E	V	A	K	V	I	V	S	M	S	I	A	F	A
AKAP-KL	P	L	E	Y	Q	A	G	L	L	V	Q	N	A	I	Q	Q	A	I
MAP-2	T	A	E	E	V	S	A	R	I	V	Q	V	V	T	A	E	A	V
EZRIN	S	Q	E	Q	L	A	A	E	L	A	E	Y	T	A	K	I	A	L
Consensus	X	(L)	X	X	X	(A)	X	X	(I)	(V)	X	X	(V)	(I)	X	X	(A)	(V)
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18



J Mol Biol. The biological functions of A-kinase anchor proteins.

Figure 3. Consensus sequence of AKAP-RII-binding domains.

Consensus sequence derived from the alignment of the primary sequences of several AKAPs and the amphipathic helical wheel and the residues forming it are depicted as a thick line.

1.4 The ubiquitin proteasome system and E3 ubiquitin ligase CHIP

In mammalian cells the post-translational modification of proteins is a common mechanism of cell regulation. The covalent modification of proteins by attachment of other protein is one such example. The control of this cellular mechanism is most important because every injury that affects this mechanism can lead to development of human diseases or disorders, including cancer. The balance between the synthesis and degradation of proteins is regulated by the ubiquitin-proteasome system (UPS).

Ubiquitylation has a central role in several physiological processes and it is involved in the regulation of cell survival, differentiation, genetic integrity, protein quality control and signaling. Frequently, the substrates of ubiquitin are degraded through the proteasome³⁷.

This process requires the activity of three enzymes: E1 (ubiquitin activating), E2 (ubiquitin conjugating) and E3 (ubiquitin ligating) that act in series to catalyze ubiquitination.

The E1 enzyme is the activating enzyme which ubiquitin is attached to in an ATP-dependent reaction. The E2 enzyme is the conjugating enzyme, which the ubiquitin is transferred to, from the E1. The E3 is the ubiquitin ligase, which directly or indirectly catalyzes the transfer of the ubiquitin to the lysine of a target protein, with the formation of an isopeptide bond³⁸ (**Figure. 4**).

The ubiquitin substrates are not always directed to degradation via UPS. Infact the amount of ubiquitin tagged protein is balanced through the activity of deubiquitylating enzymes (DUBs) that reverse ubiquitylation by removing conjugated ubiquitin tags^{39, 40}. The RING finger domain of E3 ubiquitin ligases contain a characteristic cysteine-rich-zinc-binding


domain composed by a pattern of conserved cysteine and histidine residues.

The carboxyl terminus of Hsp70-interacting protein (CHIP), also known as STUB1 (STIP1 homology and U-box containing protein1), is a member of E3 ubiquitin ligase that plays an important role in maintenance the protein homeostasis in the cytoplasm⁴¹.

In literature is described that CHIP binds several members of the molecular chaperones Hsp70/90 family that have a central role in the refolding of proteins^{42, 43}. Specifically, the principal activity of CHIP is to remove, through the ubiquitin proteasome system, the misfolded or damaged proteins that can lead to development of human cancers or other disorders^{44, 45}.

The protein CHIP was first characterized in human heart⁴². The important domain that allows the binding between CHIP and Hsp70/90 is a tetratricopeptide repeats domain (TPR) located at N-terminus of the protein, whereas a U-box domain at the C-terminus of CHIP displays the ubiquitin ligase activity.

In literature are described several ciliopathies linked to a loss activity of CHIP, in particular several form of ataxia such as spinocerebellar ataxia autosomal recessive 16 (SCAR16)^{46, 47}.



Copyright ©<http://www.progenra.com>

Figure 4. Schematic representation of ubiquitin system.

The conjugation of ubiquitin molecules to substrates requires coordinated action of three enzymes: the ubiquitin activating enzyme (E1), the ubiquitin conjugating enzyme E2 and the E3 ligase that associates the ubiquitin molecules to the substrates. Once ubiquitinated, the proteins can be degraded by the proteasome or de-ubiquitinated by a specific DUBs enzyme.

1.5 Ciliogenesis

The primary cilium is considered a sensory organelle able to receive extracellular signals and transmit them into cells.


In mammalian cells primary cilia acts as “antennae” to sense signals such as growth factors, hormones, odorants and development morphogens^{48, 49}. In the last years the focus on primary cilium has increased since this organelle has a critical role in regulating of different signaling pathways and during vertebrate development and tissue homeostasis⁵⁰.

Defects in ciliary assembly or its function can lead to several cilium-related human diseases called ciliopathies⁵¹. Ciliopathies are a group of genetic diseases caused by alterations of development, functioning and signaling of primary cilium. These syndromes present manifestations as polydactyly, retinal degeneration, mental retardation, anosmia, obesity and kidney cysts. The list of ciliopathies continues to grow and at the present includes Bardet-Biedl Syndrome (BBS), Joubert Syndrome (JS), Oral-facial-digital Type I (OFD1), retinal degeneration, polycystic kidney disease⁵².

Primary cilium usually forms during G1 phase of the cell cycle or when cells, deprived of nutrients and mitogens, exit from cell cycle and enter in quiescence state.

The primary cilium grows out from centrosome, the main microtubule-organizing center (MTOC) in animal cells, and it is composed of a microtubule-based core structure called axoneme. The axoneme is nucleated by basal body that includes the mother centriole and associated pericentriolar material (PCM). Moreover the axoneme is surrounded by a ciliary membrane and it is assembled by nine parallel doublet microtubules which elongate from the basal body. The distal region of basal body, where the outer doublets begin to form, is called the transition zone. The ciliary pocket, an invagination of

the plasma membrane at the root of the cilium, is found on some types of mammalian cells⁵³. (**Fig. 5**)



Nature Neurology. Primary cilia in neurodevelopmental disorders

Figure 5. Structure of primary cilium.

Since the protein synthesis occurs into the cytoplasm, elongation of primary cilium requires the intraflagellar-transport machinery (IFT), two complexes that move themselves within the cilium. IFT complex B is responsible of anterograde transport of ciliary proteins from base to the tip of cilium, whereas IFT complex A is responsible of retrograde transport from tip to base of cilium⁵³. In literature is described that IFT complex B is crucial for a correct assembly of primary cilia. In fact the lack of some IFT complex proteins causes short or absent cilia. By contrast,

IFT complex A transport back proteins to the basal body but it seems not be crucial for assembly of cilia⁵⁴.

In the cells there are a lot of positive and negative regulators that control the correct assembly of primary cilium. Examples of negative regulators are the centriolar protein Cp110 and the kinesin Kif24; they are able to destroy ciliary axoneme, so their abundance is lower during cilia assembly^{55, 56}. Conversely, other proteins such as the ser/Thr kinase TTBK2 and MARK4 act to promote cilium formation. Upon serum deprivation, in fact, TTBK2 localizes at basal body where removes Cp110 and recruits IFT complexes⁵⁷. The balance between cilia assembly or disassembly is regulated by also post-translational modifications. HEF1/Cas-L/NEDD9 is a component of focal adhesions that colocalizes with Aurora kinase A at the centrosome. Aurora A stimulates histone deacetylase 6(HDAC6) resulting in deacetylation of axonemal microtubules rendering them unstable⁵⁸.

1.6 Correlation between cAMP signaling, the UPS system and primary cilium

In literature is extensively documented the tight correlation between ubiquitylation and cAMP pathway while the linkage that couples the cAMP cascade and primary cilium is a very current topic.

Nevertheless several components of cAMP pathway, including G-protein coupled receptors (GPCRs), adenylate cyclases (ACs) and cAMP-dependent protein kinase A (PKA), conduct different important roles within the ciliary compartment⁵⁹.

In literature is documented that a pool of PKA is localized at centrosome, the basal structure of primary cilia⁶⁰⁻⁶².

In particular, PKA is a negative regulator of hedgehog (Hh) pathway that plays a critical role in embryonic development^{63, 64}.

In the absence of Hedgehog ligand, PKA phosphorylates Ci/Gli transcription factors promoting their proteolysis and the production of the repressor forms of Ci/Gli blocking Hedgehog target gene expression. In contrast, the activation of Hedgehog signaling increases the active forms of Ci/Gli resulting in Hedgehog target gene expression.

The basal level of PKA activity in Hedgehog-responsive cells is precisely regulated and it is maintained at the basal body of cilium by interacting with A-Kinase-Anchor-Proteins (AKAPs)⁶². Probably, this regulation is conducted by another ciliary G-coupled receptor Gpr161 that, after stimulation of the transmembrane protein Smoothened (Smo), exits from the cilium maintaining inactive the PKA⁶⁵.

This strongly suggests the existence of a localized pool of PKA maintained at the base of cilium which targets Hh signaling during the essential steps of ciliogenesis.

cAMP is a second messenger implicate in a wide of biological functions including the activity of several E3 ligases. In this manner, PKA modulating the activity of the principal enzymes

of ubiquity proteasome system, it controls the stability, the turnover and the biological activity of several cellular substrates. In neurons, PKA controls the neurite outgrowth, morphogenesis and improve the synaptic plasticity and memory. In response to an increase of intracellular levels of cAMP, the E3 ubiquitin ligase praja2 ubiquitinates and degrades NOGO-A, an important inhibitor of neurite outgrowth in mammalian brain⁶⁶.

Another one important correlation between cAMP cascade and UPS system is the regulation of the turnover of regulatory subunits of PKA by E3 ubiquitin ligase praja2. When in the cells there is an increase of cAMP levels, praja2 promote ubiquitylation and subsequent proteolysis via UPS of R subunits, regulate the strength and duration of PKA signal in response to cAMP⁶⁷.

This relationship between these two systems suggest the exists of a circuit finely regulated in which cAMP pathway controls the turnover/stability of key elements of metabolic and proliferative pathways, but at the same time UPS regulates the stability of components of the cAMP cascade and the duration and amplitude of its signal⁷.

1.7 NimA related kinase 10 (Nek10)

During the years, the scientific research has explained in which way the damages to cell cycle, checkpoint alteration and chromosome instability, can lead to development of cancers and other disorders. NIMA-related kinases (NEK) proteins are serine/threonine kinase, involved in the regulation of cell cycle, were identified in several organisms from protists to multicellular eukaryotes including mice and humans^{68, 69}. In literature is described that some members of this family are involved in ciliary functions and ciliopathies⁷⁰.

Statistical analysis have confirmed that this family of proteins coevolved with centrioles, which represent the microtubule-organizing center and prime the assembly of basal bodies of cilia⁷¹.


In human cells there are eleven genes that encode from NEK1 to NEK11 proteins. **(Fig. 6)**

Nek10 gene localizes on the short arm of human chromosome 3 (3p24.1). This gene encodes for fourteen transcripts the longest of which has 39 exons. The 4.25 Kbp transcript encodes a protein of 1172 residues with an estimate weight of 133 Kda⁷². Despite low overall sequence homology, the organizational features of NimA are broadly conserved among mammalian Nek kinases. Infact all these proteins are characterized by a N-terminal catalytic domain, except Nek10 that have its kinase domain in the central position. In addition to catalytic domain there are: His-Arg-Asp (HRD) motif which is typical of ser/thr kinases regulated through phosphorylation⁷³, coiled-coiled domains which mediate the oligomerization and PEST sequences which participate in ubiquitin dependent proteolysis⁶⁹.

In contrast with the conserved catalytic domain, the C-terminal region of NEK proteins is different in length, sequence and domain organization. As explained before, several NEK

members have important roles in cell cycle control, in particular NEK2 facilitates spindle pole separation whereas NEK6, NEK7 and NEK9 are important in generating the mitotic spindle^{74, 75}. NEK1 is involved in the repair of DNA strand breaks at G1-S and G2-M transitions⁷⁶⁻⁷⁸ and NEK10 and NEK11 are involved in G2-M DDR checkpoint.

Specially, NEK10 is required for the activation of extracellular signal-regulated kinase 1/2 (ERK1/2) signaling upon UV irradiation, but not in response to mitogens, such as epidermal growth factor. After the stimulation, NEK10 physically associated with Raf-1 and MEK1 in a Raf-1-dependent manner and the formation of this complex was necessary for Nek10-mediated MEK1 activation. The appropriate maintenance of the G2/M checkpoint following UV irradiation required Nek10 expression and ERK1/2 activation, indicating a role for Nek10 in the cellular response to UV irradiation⁶⁸.



Cell Div. Nek family of kinases in cell cycle, checkpoint control and cancer.

Fig.6 The human NIMA-related protein kinase (NEK) family

A schematic representation of human NEKs gene. Are indicated: the kinase domains (purple), coiled-coils (green), degradation motifs (red), RCC1 (regulator of chromatin condensation 1) domains (light blue) and armadillo repeats (yellow)

2 AIM OF THESIS

The cAMP signaling pathway has been carefully studied in the laboratory where I practiced my PhD program. In the last years, our attention was focused on the relationship between ciliogenesis and cAMP pathway, in particular on the regulation of the primary cilium stability, via UPS, in response to GPCR signaling. By a proteomic analysis, we identified PKA as a component of a macromolecular complex that includes the pericentriolar matrix protein 1 (PCM1) and Nima-related Kinase 10 (NEK10). PCM1 is a scaffold protein mostly localized in centriolar satellites and its role about the ciliogenesis is abundantly described⁷⁹, whereas the involvement of NEK10 kinase is widely unknown.

During my PhD program, I studied the role of NEK10 in primary ciliogenesis. I found that NEK10 plays a major role in the formation of primary cilium. Activation of GPCR-cAMP signaling causes the disassembly of primary cilium and that this stimulus primes the degradation of NEK10 protein by the ubiquitin proteasome system through E3 ubiquitin ligase CHIP. Disappearance of NEK10 levels leads to cilia resorption, underlying the central role of cAMP-NEK10 axis in the control of primary cilium stability.

Accordingly, the principal aims of my PhD thesis were the following:

1. Identify NEK10 and PKA as novel components of the multimeric signaling complex assembled at pericentriolar region by PCM1.
2. Determine the role and the mechanism of NEK10 and PKA in the control of ciliogenesis.
3. Analyze the intersection between cAMP signaling and NEK10-regulated ciliogenesis.

4. Study the role of NEK10 in the ciliopathies, such as the autosomal recessive spinocerebellar ataxia-16 (SCAR16).

3 MATERIALS AND METHODS

3.1 Cell lines. Human embryonic kidney cell line (HEK293) and primary skin fibroblasts from SCAR16 patients were cultured in DMEM containing 10% fetal bovine serum (FBS) supplemented with 2mM L-glutamine, 100 IU/ml penicillin, in an atmosphere of 5% CO₂ at 37°C.

3.2 Plasmids, siRNAs and transfection. Vectors encoding for NEK10-flag (wild type and mutants) and PCM1-HA were provided by Dr Stambolic V. and Dr Kamiya A. respectively. HA-Ub, CHIP-myc (wild type and K30A mutant), HSP70-V5, were provided by Dr Carlomagno F. and epitope Myc tagged RII β vectors were kindly provided by Dr Ginsberg SH. NEK10 phosphorylation mutants (T223A and T812A) were generated by PCR using specific oligonucleotides. siRNAs targeting distinct segments of coding regions of NEK10 and CHIP were purchased from IDT and Life technologies.

The siRNA sequence (IDT) targeting the 3'-UTR (untranslated region) of human NEK10: sense sequence: CCACAAGACAUUAGUAAA UUUACTT antisense sequence: CGGGUGUUCUGUAAUC AUUUAAAUGAA or human CHIP sense sequence: UUACACCAACCGGGCCUUt; antisense sequence: CAAGGCC GGUUGGUGUAAta.

siRNAs were transiently transfected using Lipofectamine 2000 (Invitrogen) at a final concentration of 100 pmol/ml of culture medium.

3.3 Antibodies and chemicals. Polyclonal antibodies directed against PCM1 were purchased from ABCAM and Cell Signaling and used at working 1:1000; rabbit polyclonal antibodies directed against phosphoPKA was purchased from Cell Signaling and used at working dilution of 1:1000;

monoclonal antibodies directed against RII β was purchased from BD Transduction and used at working 1:2000; haemagglutinin epitope (HA) was purchased from Covance and used at working dilution of 1:1000; monoclonal antibodies directed against flag and myc epitope used at working dilution 1:3000 were purchased from Sigma; polyclonal antibodies directed against acetylated alpha tubulin was purchased from ABCAM. Forskolin was purchased from Sigma.

3.4 Immunoprecipitation and pull down assay. Cells were washed twice with phosphate-buffered saline and lysed in a buffer (50mM TRIS–hydrogen chloride, pH 7.4, 150mM sodium chloride, 5mM magnesium chloride, 5mM dithiothreitol, 1mM ethylene diamine tetraacetic acid, 1% Triton X-100, containing aprotinin (5 μ g/ml), leupeptin (10 μ g/ml), pepstatin (2 μ g/ml), Na₃VO₄ and 1mM phenylmethylsulfonyl fluoride and protease inhibitors. The lysates were cleared by centrifugation at 15,000 g for 15 min. Cell lysates (2 mg) were immunoprecipitated in rotation at 4 °C overnight with the indicated antibodies. Pellets were washed four times in lysis buffer and eluted in Laemly buffer. An aliquot of whole cell lysates (WCE) (100 μ g) or immunoprecipitates were resolved on sodium dodecyl sulfate polyacrylamide gel and transferred on nitrocellulose membrane (Biorad, Milan, Italy) for 3 h. Filters were blocked for 1 h at room temperature in Tween-20 Phosphate buffer saline (TPBS) (PBS- Sigma, 0, 1% Tween 20, pH 7.4) containing 5% non-fat dry milk. Blots were then incubated O/N with primary antibody. Blots were washed three times with TPBS buffer and then incubated for 1 h with secondary antibody (peroxidase-coupled anti rabbit (GE-Healthcare) in TPBS. Reactive signals were revealed by enhanced ECL Western Blotting analysis system (Roche).

GST-fusions were expressed and purified from BL21 (DE3) pLysS cells. GST hybrid proteins immobilized on glutathione beads were incubated for 3 hr with cell lysates from HEK293 cells transiently expressing flag-NEK10 constructs in lysis buffer (150 mM NaCl, 50 mM Tris-HCl pH 7.5, 5mM MgCl₂, 5mM DDT, 1 mM EDTA, 1% triton X-100) in rotation at 4 °C for 4 hours. Pellets were washed four times in lysis buffer supplemented with NaCl (1 M final concentration) and eluted in Laemmli buffer. Eluted samples were size-fractionated on SDS-PAGE and immunoblotted.

3.5 PKA phosphorylation assay. Cells transfected with either wild type NEK10-flag or with NEK10-flag mutants (T223A-Flag and T812A-Flag) were left untreated or stimulated with FSK (15min). NEK10 was immunopurified with anti-flag antibodies. The precipitates were immunoblotted with anti-flag and with anti-phospho-(K/R)(K/R)X(S*/T*) specific antibodies. The quantification is shown from n=4 independent experiments (\pm SEM).

3.6 Immunofluorescence and confocal analysis. For immunofluorescence studies, HEK293 cells transiently transfected with the expression vectors were plated on poly-L-lysine (10 μ g/ml) coated glass coverslips. Cells were fixed with Paraformaldehyde for 20 minutes. After three washes, cells were immunostained with polyclonal antibodies directed against PCM1 was purchased from ABCAM, with polyclonal directed against NEK10 and used at working dilutions of 1:100; monoclonal antibodies directed against RII β was purchased from BD Transduction and used at working dilution of 1:400, monoclonal antibodies directed against Flag epitope used at working dilution of 1:400; polyclonal antibodies directed against acetylated alpha tubulin was purchased from ABCAM and used at working dilution of 1:100. High-resolution images were

acquired with a Zeiss LSM 880 confocal microscope equipped with Airyscan superresolution imaging module, using a 63×/1.40 NA Plan-Apochromat Oil DIC M27 objective lens (Zeiss MicroImaging, Jena, Germany)

3.7 Statistics Data were analyzed using analysis of variance (ANOVA) for each region and *post hoc* repeated-measure comparisons (Least Significant Difference (LSD) test). Rejection level was set at $P < 0.05$.

4 RESULTS

4.1 NEK10, RII β and PCM1 form a macromolecular complex

A proteomic analysis using full-length RII β as bait revealed that the pericentriolar matrix protein 1 (PCM1) and the Nima-related Kinase 10 (NEK10) form a complex with PKA. To verify the interaction between these proteins, I performed a co-immunoprecipitation assay (**Fig. 7a**). HEK293 cells were transfected for 24 hours with NEK10-flag vector. Lysates were immunoprecipitated with anti-RII β or non-immune IgG antibodies and the precipitates were immunoblotted with anti-PCM1, anti-flag and anti-RII β antibodies. The data in **figure 7a** show the existence of a trimeric complex composed of PKA, NEK10 and PCM1. I confirmed this interaction using an *in vitro* GST-pull down assays with the recombinant proteins (**Fig. 7b**). I transfected HEK293 cells with NEK10-flag vector and incubated the lysates with purified GST or GST-RII β fusion. The precipitates were immunoblotted with anti-PCM1, anti-flag and anti-RII β antibodies. **Figure 7b** confirmed the interaction between PCM1, NEK10 and RII β proteins.




Fig.7 (a) HEK293 cells were transiently transfected with NEK10-flag vector, lysates were immunoprecipitated with anti-RII β or non-immune IgG antibodies and immunoblotted with anti-PCM1, anti-RII β and anti-flag antibodies. **(b)** HEK293 cells were transiently transfected with NEK10-flag vector; lysates and immunoprecipitates were subjected to pull down assays with purified GST or GST-RII β fusion. The precipitates and lysates were immunoblotted with anti-PCM1, anti-flag and anti-GST antibodies.

4.2 Endogenous PCM1, NEK10 and RII β colocalize in Human Embryonic Kidney 293 cells

To demonstrate that PCM1, NEK10 and RII β are located within the same compartment, I analyzed the localization of the three proteins in HEK293 cells. For this experiment, I performed a triple immunofluorescence assay using anti-PCM1, anti-NEK10 and anti-RII β antibodies (**Fig. 8**). As shown in the figure, immunostaining analysis revealed that the three signals partially colocalize at pericentriolar region, supporting the concept that a fraction of PCM1, NEK10 and RII β proteins is restricted within the same intracellular compartment.




Fig.8 PCM1, NEK10 AND RII β colocalize in Human Embryonic Kidney cells. HEK293 were subjected to immunofluorescence assay with goat polyclonal anti-NEK10, rabbit polyclonal anti-PCM1 and mouse monoclonal anti-RII β antibodies. The merge composite of the signals shows colocalization of PCM1, NEK10 and RII β at pericentriolar region. Scale bar: 5 μ m.

4.3 NEK10 is required for ciliogenesis

NEK10 belongs to Nima related kinase family (NEKs) whose members take part to different events underlying to cell cycle progression, centrosome and microtubules formation, and mammalian ciliogenesis. Since these structures are intimately involved both in the assembly of the mitotic spindle and in ciliogenesis⁶⁹, we asked if NEK10 also contributes to the primary cilium assembly. Firstly, I demonstrated that NEK10 localizes at primary cilium. For this experiment, I deprived HEK293 cells from serum for 36 hours to induce primary cilium formation. Then, I performed a double immunofluorescence assay using anti-NEK10 and anti-acetylated tubulin antibodies. Acetylated tubulin is a modified variant of tubulin that selectively accumulates along the cilium. The immunostaining analysis shows that a significant amount of NEK10 protein localizes at the base and along the axoneme of primary cilium (Fig. 9)




Fig.9 NEK10 localizes at primary cilium. HEK293 cells were serum-deprived for 36 hours and then immunostained for NEK10 (red) and acetylated tubulin (green) antibodies. The signal was analyzed by confocal microscope equipped with Airyscan super resolution imaging module. A merge composite 2D and 3D of the signals shown that NEK10 localized along axoneme of primary cilium.

Once demonstrated the localization of NEK10 in human cells along the primary cilium, I verified the role of NEK10 during

primary cilium assembly. To this aim, I transfected transiently HEK293 cells with siRNA targeting endogenous NEK10 and deprived cells from serum for 36 hours. The data shows that, in control (siRNAc) cells, serum deprivation significantly increased the number of primary cilia. In contrast, genetic knock-down of NEK10 drastically reduced the number of ciliated cells. Re-expression of NEK10 reversed the effects of NEK10 silencing, indicating that NEK10 is, indeed, a biologically relevant player of ciliogenesis. (**Fig 10.a**). Moreover, I repeated the experiment using the kinase-dead mutant of NEK10 (NEK10-KD) carrying an inactivating mutation within the catalytic domain (K548R). As shown in the figure, the NEK10 mutant critically reduced the number of ciliated cells. In the **Fig 10.b** are showed the levels of NEK10 in siRNA transfected cells monitored by immunostaining.




Fig.10 (a) NEK10 is required for ciliogenesis. HEK293 cells were transiently transfected with control (siCNT) or with siRNAs targeting NEK10 (siNEK10), serum deprived for 36 hours, fixed and immunostained with anti-acetylated alpha tubulin and with anti-Flag antibodies (for NEK10-flag). Where indicated, NEK10-flag vector (either wild type or kinase dead, KD) was included in the siRNAs transfection mixture. Arrows indicate the localization of the cilium in cells expressing flag-tagged NEK10. Cumulative data from three independent experiments are shown (lower right panel). **(b)** The levels of NEK10 in siRNA-transfected cells were monitored by immunostaining.

4.4 PKA regulates the stability of primary cilium

The experiments above indicate that NEK10, PCM1 and PKA form a stable complex at pericentriolar region. A fraction of NEK10 localizes at- and regulates primary cilium formation. Previous work demonstrated that a pool of PKA is localized at the base of primary cilium through interaction with an as yet identified scaffold protein⁶². Localization of PKA at cilium is required for cilium formation. However, the impact of PKA activation cilium stability was largely unknown. Accordingly, I investigated the role of PKA activation on cilium stability. HEK293 cells where serum deprived for 36 hours and then stimulated with forskolin (FSK), a diterpene that activates adenylate cyclase (AC), or with isoproterenol (ISO), a beta-adrenergic receptor (bAR) agonist. Cells were subjected to immunostaining analysis using anti-acetylated tubulin antibody. As shown in **Fig.11**, treatment with FSK or Isoprotenerol strongly decreased the number of ciliated cells. These data suggested that activation of cAMP signaling critically impacts on primary cilium stability.




Fig.11 cAMP induced resorption of primary cilium. HEK293 cells were serum deprived for 36h and then left untreated (CNT) or stimulated with isoproterenol (Iso) or forskolin (FSK) for 3 hours. The same cells were immunostained with anti-acetylated tubulin and Draq5. Cumulative data from five independent experiments are shown (lower right panel)

4.5 PKA phosphorylation primes NEK10 for proteolysis via UPS

Our data indicate that NEK10 is required for ciliogenesis, while PKA activation induces cilium resorption. Since NEK10 and PKA are present within the same multimeric complex assembled by PCM1, I tested if/how PKA activation regulates NEK10 levels. To this aim, I monitored the levels of NEK10 in HEK293 cells after the treatment with FSK for 1 hour. The data shown in **Fig.12** revealed that the activation of PKA by FSK caused a severe decrease of NEK10 levels. The effects of forskolin on NEK10 levels were abrogated by pre-treating the cells with the proteasome inhibitor MG132.

The data indicate that, in response to cAMP stimulation, NEK10 undergoes to proteasomal degradation.




Fig.12 PKA stimulation induces proteolysis of NEK10 by the proteasome. HEK293 cells were transiently transfected with NEK10-flag. After 24 hours, cells were harvested and treated with forskolin (40 μ M). Where indicated, cells were pretreated with MG132 (20 μ M/3 hours). Lysates were immunoblotted with anti-flag or anti-tubulin antibodies.

The data above indicate that, in response to cAMP stimulation, NEK10 undergoes proteasomal degradation. We assume that PKA phosphorylation primes NEK10 for proteolysis. Primary sequence analysis of NEK10 predicts two conserved PKA phosphorylation sites (T223 and T812) (**Fig.13**).




Fig.13 Schematic diagram showing the protein sequence of human NEK10 and the putative PKA consensus sites (thr223 and thr812).

To ask if phosphorylation of one or both of these sites renders NEK10 susceptible to proteolysis, we generated mutant forms of NEK10 using a site-directed mutagenesis to substitute either T223 or T812 with alanine. We tested our hypothesis by analyzing the phosphorylation status of affinity-isolated NEK10 with a PKA substrate antibody. In contrast to phosphorylation of the wild type and T223A NEK10 mutant, the substitution of T812A abolished both basal and FSK-induced NEK10 phosphorylation (**Fig.14**).




Fig.14 PKA phosphorylates NEK10 at threonine 812. HEK293 cells were transfected with either wild type NEK10-flag or with NEK10-flag mutants (T223A-Flag and T812A-Flag) and were left untreated or stimulated with FSK (40 μ M 715 min). NEK10 was immunopurified with anti-flag antibodies and the precipitates were immunoblotted with anti-flag and anti-phospho-(K/R) (K/R) X(S*/T*) specific antibodies.

Next, we asked if this phosphorylation by PKA is required to induce NEK10 proteolysis. To verify this hypothesis, I evaluated the levels of NEK10 in cells transfected either with NEK10 wild type or with T812A mutant vectors. Both cell lines were stimulated with FSK (**Fig.15a**) or Isoproterenol (**Fig.15b**) for 30 and 60 minutes. The figures show that the phospho mutant (T812A), not sensible to phosphorylation by PKA, is not degraded by FSK and ISO compared to wild type protein.

Ubiquitination is required for proteasomal degradation of a variety of cellular substrates⁸⁰. Accordingly, I asked if cAMP induces ubiquitination of NEK10. To test this hypothesis, I performed ubiquitination assays in HEK293 cells transfected either with hemagglutinin (HA)-tagged ubiquitin and NEK10 (wild type or T812A mutant) and treated with FSK for 60 minutes. The lysates were immunoprecipitates with anti-flag

antibody and the precipitates were immunoblotted with anti-HA antibody. **Fig.15c** shows that FSK induces the accumulation of poly-ubiquitinated forms of NEK10, whereas this poly-ubiquitination was abrogated by the T812A mutation. These experiments confirmed that phosphorylation by PKA is necessary to prime proteolysis of NEK10 via UPS.




Fig.15 (a) Phosphorylation by PKA drives proteolysis of NEK10. (a-b) Cells transfected with either wild type NEK10-flag or with T812A-flag mutant were left untreated or stimulated with forskolin (**a**) or isoproterenol (**b**) for 30-60 min. Total cell lysates were immunoblotted with anti-flag and anti-tubulin antibodies. (**c**) Cells were transiently co-transfected with NEK10-flag construct (either wild type or T812A mutant) and HA-ubiquitin. Lysates were subjected to immunoprecipitation with anti-flag and immunoblotted with anti-HA and anti-flag antibodies.

Next, I tested if NEK10 phosphorylation was required for primary cilium disassembly induced by the cAMP cascade. Cells were transiently transfected with NEK10 (either wild type or the NEK10-T812A mutant), serum-deprived for two days and then treated with FSK. As shown in **Fig.16**, the T812A mutation prevented FSK-induced cilia disassembly, supporting the concept that PKA phosphorylation of T812 primes NEK10 for proteolysis, which results in cilia disassembly.




Fig.16. Phosphorylation of NEK10 by PKA induced cilia resorption. HEK293 cells were transfected with either wild type NEK10-flag or with T812A-flag mutant. After transfection, cells were serum deprived for 36h and left untreated or stimulated with FSK (40 μ M/3 hours). Cells were subjected to a double immunofluorescence for flag (red) and acetylated tubulin (green). Cumulative data from three independent experiments are shown on right panel.

4.6 CHIP is the NEK10 E3 ubiquitin ligase

Since NEK10 is efficiently ubiquitinated under cAMP stimulation, it was necessary to identify the E3 ligase responsible for this ubiquitination. A proteomic analysis identified several PKA partners; one of this is the E3 ubiquitin ligase C-terminus of HSP70 interacting protein (CHIP) known as STUB1. CHIP is a chaperone-associated E3ligase involved in the ubiquitination and degradation of HSP70-bound substrates and contains a tetra-tricopeptide (TPR) motif tandem repeats that mediates interaction with HSP70. First, I verified the interaction between the two proteins. I performed a co-immunoprecipitation assay using lysates from cells transfected with NEK10-flag, HSP70-V5 and CHIP-Myc vectors. The lysates were immunoprecipitated with anti-myc antibody and precipitates were immunoblotted with anti-flag, anti-V5 and anti-myc antibodies. As shown in **Fig.17a** the three proteins form a stable complex in cell lysate.

Then, I tested if the binding between NEK10, CHIP and HSP70 was regulated by cAMP. To this aim, I performed a co-immunoprecipitation assay using lysates from cells transfected either with NEK10 wild type or with NEK-T812A mutant. Cells were induced with FSK for 30 minutes; the lysates were immunoprecipitated with anti-myc antibody and immunoblotted with anti-flag, anti-V5 and anti-myc antibodies. The **Fig.17b-c** shows that the binding between NEK10, HSP70 and CHIP increases after the treatment with FSK, in contrast, the T812A mutation significantly decreases NEK10 binding to CHIP.




Fig.17 (a) cAMP-induced binding of CHIP to NEK10. HEK293 cells were transiently co-transfected with NEK10-flag, HSP70-V5 and CHIP-myc vectors. To prevent NEK10 degradation by CHIP, cells were treated with MG132 (20 μ M/8hours) before harvesting. Lysates were immunoprecipitated with anti-flag or with control IgG. The precipitates and lysates were immunoblotted with the indicated antibodies. **(b)** Cells were co-transfected with NEK10-flag vectors (either wild type or T812A mutant), HSP70-V5 and CHIP-myc, serum deprived for 24 hours and then left untreated or stimulated with FSK (40 μ M/30 min). Lysates were immunoprecipitated with anti-flag antibody and were immunoblotted with the indicated antibodies. **(c)** Cumulative data of three independent experiments shown in b.

We then asked if CHIP degrades NEK10 in the absence of MG132. In the **Fig. 17d** I co-transfected cells with NEK10-flag and CHIP-myc vectors, alternately with wild type or with catalytically inactive mutant (K30A) of CHIP that does not bind HSP70. The lysates were immunoblotted with anti flag, myc and tubulin antibodies. The figure shows that in presence of CHIP wild type there is a decrease of NEK10 levels whereas CHIP-K30A mutant in not able to degradate NEK10.

In the **Fig17.e** I performed an ubiquitination assay using cells transfected with control siRNAs or siRNAs targeting endogenous CHIP. After transfection, cells were left untreated or stimulated with Isoprotenerol for 1 hour. The figure shows that in basal condition there is an accumulation of poly-ubiquitinated form of NEK10 ISO dependent, by the contrast, the genetic knock-down of endogenous CHIP prevented ISO-induced NEK10 polyubiquitination. These findings supported the idea that PKA controls NEK10 stability through CHIP.




Fig.17 CHIP ubiquitylates NEK10. (d) Lysates from cells co-transfected with NEK10-flag and CHIP (either wild type or K30A mutant) were immunoblotted with the indicated antibodies (e) Cells co-transfected with HA-ubiquitin, NEK10-flag and siRNAs (either control siRNA or siCHIP) were serum-deprived overnight and stimulated with isoproterenol for 1 hour. Lysates were subjected to immunoprecipitation with anti-flag antibody. Ubiquitinated NEK10 was revealed by immunoblot with anti-HA antibodies.

Once determined that cAMP primes proteolysis of NEK10 through the interaction with CHIP, we verified if CHIP mediates the effects of cAMP on cilia stability. To this aim, I transfected HEK293 cells with control siRNAs or siRNAs targeting endogenous CHIP. Twenty hours after the transfection I deprived the cells from serum for 36 hours and treated them with FSK for 3 hours. Cells were subjected to immunofluorescence assay with anti-acetylated tubulin antibody. As shown in the figure (**Fig 17f**), down regulation of endogenous CHIP prevented cilia resorption induced by FSK treatment.




Fig.17 CHIP ubiquitylates NEK10 and mediates cAMP effects on primary cilium stability (f) Cells were transiently transfected with control or with siRNAs targeting CHIP, serum-deprived for 36h and then left untreated or stimulated with FSK (40 μ M) for 3 hours. Primary cilia were visualized by immunostaining with acetylated tubulin antibody whereas nuclei with Draq5. Cumulative data from five independent experiments are shown in the graph near the figure.

4.7 Dysregulation of CHIP affects cilia in SCAR16 disease

Biallelic *STUB1* mutations resulting in aberrant CHIP have been identified in patients with clinical features of autosomal recessive spinocerebellar ataxia-16 (SCAR16). This is a rare genetic syndrome characterized by truncal and limb ataxia resulting in gait instability, mild peripheral sensory neuropathy, and cognitive defects. Hypogonadism can also be present in these patients (Gordon Holmes syndrome, GHS), consistent with signaling defects and altered responses to hypothalamic hormones. Mice lacking *STUB1*/CHIP gene show a phenotype that recapitulates most of the SCAR16 features⁸¹. Accordingly, we determined if CHIP mutations affect primary cilia. We analyzed ciliogenesis in primary fibroblasts isolated from cutaneous biopsies of SCAR16 patients or from healthy volunteers. **Fig.18** shows that FSK treatment in normal fibroblasts promoted resorption of cilia. In contrast, no major effects of FSK stimulation on cilia were evident in SCAR16 fibroblasts. Interestingly, genetic silencing of *NEK10* in SCAR16 fibroblasts markedly reduced the number of ciliated cells, even in the absence of FSK, further supporting a role of the CHIP-*NEK10* axis in control of cilium stability.




Fig.18 CHIP, NEK10 and cilia in SCAR16 fibroblasts. Skin fibroblasts from healthy volunteers (BJ) and SCAR16 patients (AX71) were serum deprived for 48h and treated with FSK (80 μM/6h). Cells were fixed and stained for acetylated tubulin and Draq5. Where indicated, AX71 cells were transiently transfected with control siRNA or with siRNA targeting endogenous NEK10, before stimulation. Cumulative data from 4 independent experiments are shown.

5 DISCUSSION AND CONCLUSION

Primary cilia are considered very important organelles that emanate from cell surface and are able to detect external signals and reintroduce them into cells. Primary cilia are present on the apical surface of the majority of cells in the human body and the structure that promotes the assembly of primary cilium is the centrosome, the principal microtubule organizing center (MTOC) in animal cells. The cilium is composed by axoneme and the basal body⁵³.

The primary cilium is a compartmentalized complex for signal integration and propagation relevant for many developmental processes. In dividing cells, the transition between centrosome and primary cilium is functionally linked. In mitotic interphase, centrosomes organize the cytoplasmic microtubule network, whereas in mitosis they regulate mitotic spindle dynamics and cytokinesis. In postmitotic cells, the centrosome migrates to the cell surface, and one of the centrioles differentiates into a basal body from which microtubules nucleate to form a primary cilium. In normal proliferating cells, the cilium can be transiently observed in G1 phase, disappearing when the cell enters the cell cycle⁸². A significant fraction of PKA is localized at the base of cilium through interaction with AKAPs, controlling essential aspects of ciliogenesis and the Hedgehog (Hh) pathway. However, the impact of PKA activation on the turnover of ciliary proteins and its role in primary cilium stability were largely unknown.

As mentioned above, there is a tight connection between cAMP cascade and UPS system and derangements in these mechanisms are linked to arise several neurodegenerative and proliferative disorders.

In the thesis, I reported the identification of the serine/threonine kinase NEK10 as a novel positive regulator of ciliogenesis. NEK10 is member of the Nima-related kinases activated at

G2/M transition and its activity is required for proper cell cycle progression. I demonstrated the existence of a macromolecular complex in which NEK10 is localized at the primary cilium through the interaction with PKA and PCM1, a pericentriolar scaffold protein involved in different aspects of microtubule dynamics, cell division and ciliogenesis. I found that NEK10 protein stability is a critical determinant for the assembly/disassembly of cilium and it is mediated by the GPCR signaling. By a combinatorial approach of biochemistry, cell biology and molecular genetics, I demonstrate the presence of a trimeric complex composed by NEK10, PCM1 and regulatory subunit of PKA (RII β) at pericentriolar region of mammalian cells. In particular, I found that NEK10 localizes at the base and along the axoneme of primary cilium. NEK10 down regulation severely affected the assembly of primary cilium.

I also analyzed the intersection between GPCR signaling and primary cilium. I found that NEK10 is a novel direct target of PKA. Phosphorylation of NEK10 by PKA at Thr812 primes NEK10 to ubiquitination and proteolysis. Disappearance of NEK10 promotes cilia resorption. A proteomic analysis allowed the identification of CHIP as the E3 ubiquitin ligase that binds to- and ubiquitylates NEK10, causing NEK10 proteolysis through the UPS and cilia resorption. Removal of CHIP prevented cAMP effects on cilium resorption. These findings point to CHIP as a novel regulator of protein turnover at ciliary sites that efficiently couples GPCR signaling to cilia dynamics. This mode of regulation was further supported by evidence that germline inactivating mutations of CHIP that cause SCAR16 disease prevented cAMP-induced disassembly of cilia.

Altogether, the findings reported in my thesis elucidate the mechanism(s) underlying cilia resorption during GPCR stimulation, both in healthy and disease conditions. They also provide mechanistic insights into how cAMP controls cell growth. It is well established that the cAMP cascade regulates

growth and differentiation of a wide variety of cell types. PKA activation can either induce or inhibit cell growth, depending on cell type or metabolic conditions⁸³. In growth-arrested endocrine cells, the cAMP-PKA pathway promotes the transition from G0 to G1 phase, allowing the cells to progress through the cell cycle⁸⁴. The transition from quiescent to proliferative state requires disassembly of the primary cilium.

Several targets of PKA have been identified and causally linked to induction of cell growth. However, if and how PKA activation modulates the activity of proteins controlling cilia stability in starved cells was largely unexplored. These findings help to define the relevance of PKA pathway in cilia resorption in the course of hormone stimulation. We show that PKA activation by cAMP agonists targets NEK10 for proteolysis through the UPS. The cAMP cascade induces cilia disassembly and promotes entry into the cell cycle by removing the NEK10 pro-ciliogenic kinase. NEK10 thus represents a nodal point in the ciliary compartment where cAMP signaling and the UPS converge and integrate to control essential aspects of cilia dynamics and, most likely, cell growth. Mutations affecting any component of this proteolytic machinery may alter the sensitivity of the cells to hormones or growth factors, profoundly impacting on cell growth and vertebrate development.

Although the results of my thesis enhance the role of cAMP into disassembly of primary cilium, there are some points that need to be addressed. It is important to understand whether cilia resorption induced by cAMP has a physiologically relevant implications for cell biology. Previous work revealed that activation of cAMP pathway promotes ciliogenesis.^{85, 86}

This apparent discrepancy could not be ascribed to the different cell models used, since we confirmed that in serum supplemented, confluent cells cAMP stimulation had no major impact on cilium stability (data not shown). These findings

suggest that cAMP pathway may have a dual effect on primary cilia depending on how growth arrest is achieved. In the presence of serum, cAMP contributes to primary ciliogenesis induced by cell confluency, while under serum starvation the same messenger promotes cilium disassembly.

Finally, the findings reported in my thesis indicate that NEK10 is a new ciliary protein that in concert with other centriolar proteins plays a major role in the regulation of assembly/disassembly of primary cilium. My next goal is the identification and the molecular characterization of the relevant NEK10 substrates involved in mammalian ciliogenesis. Derangement of this NEK10-regulated signaling circuitry may underpin to genetic and proliferative disorders human disorders.

In conclusion, I have identified a PCM1-centered multimeric complex that functionally links second messenger signaling (cAMP), kinase activities (PKA, NEK10) and the UPS (CHIP) to cilia dynamics. This mechanism explains how compartmentalized signaling networks regulate cilia formation in both physiological and pathological conditions.




Figure 19. Molecular mechanism model. Under resting conditions, PKA holoenzyme form a stable complex with PCM1 and NEK10, promoting ciliogenesis. Elevation of intracellular cAMP levels by ligand (L) stimulation of the adenylate cyclase (AC) efficiently activates PKA which phosphorylates NEK10. Phosphorylation primes ubiquitin-dependent proteolysis of NEK10 by E3 ligase CHIP. The decrease of NEK10 levels promotes the resorption of primary cilium.

6 APPENDICES

AC Adenylyl cyclase

GPCR G protein coupled receptor

PKA Protein kinase A

PDE Phosphodiesterase

UPS Ubiquitin Proteasome System

CHIP C-terminus of HSC70 interacting protein

FSK Forskolin

ISO Isoprotenerol

cAMP cyclic AMP

EPAC RAP exchange proteins

cNGC cAMP gated ion channels

AKAP A-Kinase-Anchor-Proteins

TPR tetratricopeptide repeats domain

SCAR16 spinocerebellar ataxia autosomal recessive 16

PCM1 pericentriolar matrix protein 1

NEK10 Nima-related Kinase 10

R11 β 51 regulatory subunit of PKA

Hh Hedgehog

7 ACKNOWLEDGEMENTS

For contribution in my thesis, I wish to thank Drs Eduard Stefan and Omar Torres-Quesada for their help in NEK10 phosphorylation assay and for proteomic analyses.

I would thank Drs Ivan Conte and Daniela Intartaglia for help in immunofluorescence assays with Airyscan superresolution imaging module, and Dr. Synofzik for kindly providing the fibroblast lines. Thanks to Drs Stambolic V., Kamiya A., Lim C. And Carlomagno F. For kindly sharing the vectors.

8 REFERENCES

1. Sutherland, E.W. On the biological role of cyclic AMP. *Jama* **214**, 1281-1288 (1970).
2. Taylor, S.S. *et al.* Signaling through cAMP and cAMP-dependent protein kinase: diverse strategies for drug design. *Biochimica et biophysica acta* **1784**, 16-26 (2008).
3. Rosenbaum, D.M., Rasmussen, S.G. & Kobilka, B.K. The structure and function of G-protein-coupled receptors. *Nature* **459**, 356-363 (2009).
4. Chen, J. & Iyengar, R. Inhibition of cloned adenylyl cyclases by mutant-activated Gi-alpha and specific suppression of type 2 adenylyl cyclase inhibition by phorbol ester treatment. *The Journal of biological chemistry* **268**, 12253-12256 (1993).
5. Cooper, D.M. & Tabbasum, V.G. Adenylate cyclase-centred microdomains. *The Biochemical journal* **462**, 199-213 (2014).
6. Maurice, D.H. *et al.* Advances in targeting cyclic nucleotide phosphodiesterases. *Nature reviews. Drug discovery* **13**, 290-314 (2014).
7. Rinaldi, L., Sepe, M., Donne, R.D. & Feliciello, A. A dynamic interface between ubiquitylation and cAMP signaling. *Frontiers in pharmacology* **6**, 177 (2015).
8. Kenan, Y., Murata, T., Shakur, Y., Degerman, E. & Manganiello, V.C. Functions of the N-terminal region of cyclic nucleotide phosphodiesterase 3 (PDE 3) isoforms. *The Journal of biological chemistry* **275**, 12331-12338 (2000).
9. Lomas, O. & Zaccolo, M. Phosphodiesterases maintain signaling fidelity via compartmentalization of cyclic nucleotides. *Physiology* **29**, 141-149 (2014).
10. Kim, C., Cheng, C.Y., Saldanha, S.A. & Taylor, S.S. PKA-I holoenzyme structure reveals a mechanism for cAMP-dependent activation. *Cell* **130**, 1032-1043 (2007).
11. Knighton, D.R. *et al.* Crystal structure of the catalytic subunit of cyclic adenosine monophosphate-dependent protein kinase. *Science* **253**, 407-414 (1991).
12. Taylor, S.S., Knighton, D.R., Zheng, J., Ten Eyck, L.F. & Sowadski, J.M. Structural framework for the protein kinase family. *Annual review of cell biology* **8**, 429-462 (1992).
13. Zheng, J. *et al.* 2.2 A refined crystal structure of the catalytic subunit of cAMP-dependent protein kinase complexed with MnATP and a peptide inhibitor. *Acta crystallographica. Section D, Biological crystallography* **49**, 362-365 (1993).
14. Taylor, S.S. *et al.* Dynamics of signaling by PKA. *Biochimica et biophysica acta* **1754**, 25-37 (2005).
15. Tasken, K. & Aandahl, E.M. Localized effects of cAMP mediated by distinct routes of protein kinase A. *Physiological reviews* **84**, 137-167 (2004).
16. Edelman, A.M., Blumenthal, D.K. & Krebs, E.G. Protein serine/threonine kinases. *Annual review of biochemistry* **56**, 567-613 (1987).
17. Qi, M. *et al.* Impaired hippocampal plasticity in mice lacking the Cbeta1 catalytic subunit of cAMP-dependent protein kinase. *Proceedings of the National Academy of Sciences of the United States of America* **93**, 1571-1576 (1996).
18. Brandon, E.P., Idzerda, R.L. & McKnight, G.S. Targeting the mouse genome: a compendium of knockouts (Part II). *Current biology : CB* **5**, 758-765 (1995).
19. Amieux, P.S. *et al.* Compensatory regulation of R1alpha protein levels in protein kinase A mutant mice. *The Journal of biological chemistry* **272**, 3993-3998 (1997).
20. Burton, K.A. *et al.* Type II regulatory subunits are not required for the anchoring-dependent modulation of Ca²⁺ channel activity by cAMP-dependent protein kinase. *Proceedings of the National Academy of Sciences of the United States of America* **94**, 11067-11072 (1997).
21. Burton, K.A., Treash-Osio, B., Muller, C.H., Dunphy, E.L. & McKnight, G.S. Deletion of type IIalpha regulatory subunit delocalizes protein kinase A in mouse sperm without affecting motility or fertilization. *The Journal of biological chemistry* **274**, 24131-24136 (1999).

22. Cummings, D.E. *et al.* Genetically lean mice result from targeted disruption of the RII beta subunit of protein kinase A. *Nature* **382**, 622-626 (1996).
23. McKnight, G.S. *et al.* Cyclic AMP, PKA, and the physiological regulation of adiposity. *Recent progress in hormone research* **53**, 139-159; discussion 160-131 (1998).
24. Adams, M.R. *et al.* Loss of haloperidol induced gene expression and catalepsy in protein kinase A-deficient mice. *Proceedings of the National Academy of Sciences of the United States of America* **94**, 12157-12161 (1997).
25. Brandon, E.P. *et al.* Defective motor behavior and neural gene expression in RIIbeta-protein kinase A mutant mice. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **18**, 3639-3649 (1998).
26. Fimia, G.M. & Sassone-Corsi, P. Cyclic AMP signalling. *Journal of cell science* **114**, 1971-1972 (2001).
27. Rubin, C.S. A kinase anchor proteins and the intracellular targeting of signals carried by cyclic AMP. *Biochimica et biophysica acta* **1224**, 467-479 (1994).
28. Edwards, A.S. & Scott, J.D. A-kinase anchoring proteins: protein kinase A and beyond. *Current opinion in cell biology* **12**, 217-221 (2000).
29. Dodge, K. & Scott, J.D. AKAP79 and the evolution of the AKAP model. *FEBS letters* **476**, 58-61 (2000).
30. Carr, D.W. *et al.* Interaction of the regulatory subunit (RII) of cAMP-dependent protein kinase with RII-anchoring proteins occurs through an amphipathic helix binding motif. *The Journal of biological chemistry* **266**, 14188-14192 (1991).
31. Newlon, M.G., Roy, M., Hausken, Z.E., Scott, J.D. & Jennings, P.A. The A-kinase anchoring domain of type IIalpha cAMP-dependent protein kinase is highly helical. *The Journal of biological chemistry* **272**, 23637-23644 (1997).
32. Herberg, F.W., Maleszka, A., Eide, T., Vossebein, L. & Tasken, K. Analysis of A-kinase anchoring protein (AKAP) interaction with protein kinase A (PKA) regulatory subunits: PKA isoform specificity in AKAP binding. *Journal of molecular biology* **298**, 329-339 (2000).
33. Feliciello, A., Rubin, C.S., Avvedimento, E.V. & Gottesman, M.E. Expression of a kinase anchor protein 121 is regulated by hormones in thyroid and testicular germ cells. *The Journal of biological chemistry* **273**, 23361-23366 (1998).
34. Feliciello, A., Gottesman, M.E. & Avvedimento, E.V. The biological functions of A-kinase anchor proteins. *Journal of molecular biology* **308**, 99-114 (2001).
35. Dell'Acqua, M.L. *et al.* Regulation of neuronal PKA signaling through AKAP targeting dynamics. *European journal of cell biology* **85**, 627-633 (2006).
36. Welch, E.J., Jones, B.W. & Scott, J.D. Networking with AKAPs: context-dependent regulation of anchored enzymes. *Molecular interventions* **10**, 86-97 (2010).
37. Ciechanover, A. Proteolysis: from the lysosome to ubiquitin and the proteasome. *Nature reviews. Molecular cell biology* **6**, 79-87 (2005).
38. Dikic, I. & Robertson, M. Ubiquitin ligases and beyond. *BMC biology* **10**, 22 (2012).
39. Wilkinson, K.D. DUBs at a glance. *Journal of cell science* **122**, 2325-2329 (2009).
40. Komander, D., Clague, M.J. & Urbe, S. Breaking the chains: structure and function of the deubiquitinases. *Nature reviews. Molecular cell biology* **10**, 550-563 (2009).
41. Sun, C. *et al.* Diverse roles of C-terminal Hsp70-interacting protein (CHIP) in tumorigenesis. *Journal of cancer research and clinical oncology* **140**, 189-197 (2014).
42. Ballinger, C.A. *et al.* Identification of CHIP, a novel tetratricopeptide repeat-containing protein that interacts with heat shock proteins and negatively regulates chaperone functions. *Molecular and cellular biology* **19**, 4535-4545 (1999).
43. Connell, P. *et al.* The co-chaperone CHIP regulates protein triage decisions mediated by heat-shock proteins. *Nature cell biology* **3**, 93-96 (2001).
44. Imai, Y. *et al.* CHIP is associated with Parkin, a gene responsible for familial Parkinson's disease, and enhances its ubiquitin ligase activity. *Molecular cell* **10**, 55-67 (2002).

45. Shimura, H., Schwartz, D., Gygi, S.P. & Kosik, K.S. CHIP-Hsc70 complex ubiquitinates phosphorylated tau and enhances cell survival. *The Journal of biological chemistry* **279**, 4869-4876 (2004).
46. Bettencourt, C. *et al.* Clinical and Neuropathological Features of Spastic Ataxia in a Spanish Family with Novel Compound Heterozygous Mutations in STUB1. *Cerebellum* **14**, 378-381 (2015).
47. Shi, C.H. *et al.* Ataxia and hypogonadism caused by the loss of ubiquitin ligase activity of the U box protein CHIP. *Human molecular genetics* **23**, 1013-1024 (2014).
48. Singla, V. & Reiter, J.F. The primary cilium as the cell's antenna: signaling at a sensory organelle. *Science* **313**, 629-633 (2006).
49. Barbari, N.F., O'Connor, A.K., Haycraft, C.J. & Yoder, B.K. The primary cilium as a complex signaling center. *Current biology : CB* **19**, R526-535 (2009).
50. Oh, E.C., Vasanth, S. & Katsanis, N. Metabolic regulation and energy homeostasis through the primary Cilium. *Cell metabolism* **21**, 21-31 (2015).
51. Guemez-Gamboa, A., Coufal, N.G. & Gleeson, J.G. Primary cilia in the developing and mature brain. *Neuron* **82**, 511-521 (2014).
52. Tobin, J.L. & Beales, P.L. The nonmotile ciliopathies. *Genetics in medicine : official journal of the American College of Medical Genetics* **11**, 386-402 (2009).
53. Ishikawa, H. & Marshall, W.F. Ciliogenesis: building the cell's antenna. *Nature reviews. Molecular cell biology* **12**, 222-234 (2011).
54. Marshall, W.F. & Rosenbaum, J.L. Intraflagellar transport balances continuous turnover of outer doublet microtubules: implications for flagellar length control. *The Journal of cell biology* **155**, 405-414 (2001).
55. Spektor, A., Tsang, W.Y., Khoo, D. & Dynlacht, B.D. Cep97 and CP110 suppress a cilia assembly program. *Cell* **130**, 678-690 (2007).
56. Pearson, C.G. A kinesin in command of primary ciliogenesis. *Cell* **145**, 817-819 (2011).
57. Goetz, S.C., Liem, K.F., Jr. & Anderson, K.V. The spinocerebellar ataxia-associated gene Tau tubulin kinase 2 controls the initiation of ciliogenesis. *Cell* **151**, 847-858 (2012).
58. Pugacheva, E.N., Jablonski, S.A., Hartman, T.R., Henske, E.P. & Golemis, E.A. HEF1-dependent Aurora A activation induces disassembly of the primary cilium. *Cell* **129**, 1351-1363 (2007).
59. Hilgendorf, K.I., Johnson, C.T. & Jackson, P.K. The primary cilium as a cellular receiver: organizing ciliary GPCR signaling. *Current opinion in cell biology* **39**, 84-92 (2016).
60. Nigg, E.A., Schafer, G., Hilz, H. & Eppenberger, H.M. Cyclic-AMP-dependent protein kinase type II is associated with the Golgi complex and with centrosomes. *Cell* **41**, 1039-1051 (1985).
61. De Camilli, P., Moretti, M., Donini, S.D., Walter, U. & Lohmann, S.M. Heterogeneous distribution of the cAMP receptor protein RII in the nervous system: evidence for its intracellular accumulation on microtubules, microtubule-organizing centers, and in the area of the Golgi complex. *The Journal of cell biology* **103**, 189-203 (1986).
62. Barzi, M., Berenguer, J., Menendez, A., Alvarez-Rodriguez, R. & Pons, S. Sonic-hedgehog-mediated proliferation requires the localization of PKA to the cilium base. *Journal of cell science* **123**, 62-69 (2010).
63. Matus, D.Q., Magie, C.R., Pang, K., Martindale, M.Q. & Thomsen, G.H. The Hedgehog gene family of the cnidarian, *Nematostella vectensis*, and implications for understanding metazoan Hedgehog pathway evolution. *Developmental biology* **313**, 501-518 (2008).
64. Li, W., Ohlmeyer, J.T., Lane, M.E. & Kalderon, D. Function of protein kinase A in hedgehog signal transduction and *Drosophila* imaginal disc development. *Cell* **80**, 553-562 (1995).
65. Mukhopadhyay, S. *et al.* The ciliary G-protein-coupled receptor Gpr161 negatively regulates the Sonic hedgehog pathway via cAMP signaling. *Cell* **152**, 210-223 (2013).
66. Sepe, M. *et al.* Proteolytic control of neurite outgrowth inhibitor NOGO-A by the cAMP/PKA pathway. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 15729-15734 (2014).

67. Lignitto, L. *et al.* Control of PKA stability and signalling by the RING ligase praja2. *Nature cell biology* **13**, 412-422 (2011).
68. Moniz, L.S. & Stambolic, V. Nek10 mediates G2/M cell cycle arrest and MEK autoactivation in response to UV irradiation. *Molecular and cellular biology* **31**, 30-42 (2011).
69. Fry, A.M., O'Regan, L., Sabir, S.R. & Bayliss, R. Cell cycle regulation by the NEK family of protein kinases. *Journal of cell science* **125**, 4423-4433 (2012).
70. Bettencourt-Dias, M., Hildebrandt, F., Pellman, D., Woods, G. & Godinho, S.A. Centrosomes and cilia in human disease. *Trends in genetics : TIG* **27**, 307-315 (2011).
71. Quarmbay, L.M. & Mahjoub, M.R. Caught Nek-ing: cilia and centrioles. *Journal of cell science* **118**, 5161-5169 (2005).
72. Ahmed, S. *et al.* Newly discovered breast cancer susceptibility loci on 3p24 and 17q23.2. *Nature genetics* **41**, 585-590 (2009).
73. Johnson, L.N., Noble, M.E. & Owen, D.J. Active and inactive protein kinases: structural basis for regulation. *Cell* **85**, 149-158 (1996).
74. O'Regan, L., Blot, J. & Fry, A.M. Mitotic regulation by NIMA-related kinases. *Cell division* **2**, 25 (2007).
75. Sdelci, S., Bertran, M.T. & Roig, J. Nek9, Nek6, Nek7 and the separation of centrosomes. *Cell cycle* **10**, 3816-3817 (2011).
76. Chen, Y., Chen, C.F., Riley, D.J. & Chen, P.L. Nek1 kinase functions in DNA damage response and checkpoint control through a pathway independent of ATM and ATR. *Cell cycle* **10**, 655-663 (2011).
77. Pelegrini, A.L. *et al.* Nek1 silencing slows down DNA repair and blocks DNA damage-induced cell cycle arrest. *Mutagenesis* **25**, 447-454 (2010).
78. Polci, R., Peng, A., Chen, P.L., Riley, D.J. & Chen, Y. NIMA-related protein kinase 1 is involved early in the ionizing radiation-induced DNA damage response. *Cancer research* **64**, 8800-8803 (2004).
79. Wang, G. *et al.* PCM1 recruits Plk1 to the pericentriolar matrix to promote primary cilia disassembly before mitotic entry. *Journal of cell science* **126**, 1355-1365 (2013).
80. Amm, I., Sommer, T. & Wolf, D.H. Protein quality control and elimination of protein waste: the role of the ubiquitin-proteasome system. *Biochimica et biophysica acta* **1843**, 182-196 (2014).
81. Ko, H.S. *et al.* CHIP regulates leucine-rich repeat kinase-2 ubiquitination, degradation, and toxicity. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 2897-2902 (2009).
82. Plotnikova, O.V., Golemis, E.A. & Pugacheva, E.N. Cell cycle-dependent ciliogenesis and cancer. *Cancer research* **68**, 2058-2061 (2008).
83. Schmitt, J.M. & Stork, P.J. PKA phosphorylation of Src mediates cAMP's inhibition of cell growth via Rap1. *Molecular cell* **9**, 85-94 (2002).
84. Richards, J.S. New signaling pathways for hormones and cyclic adenosine 3',5'-monophosphate action in endocrine cells. *Molecular endocrinology* **15**, 209-218 (2001).
85. Ou, Y. *et al.* Adenylate cyclase regulates elongation of mammalian primary cilia. *Experimental cell research* **315**, 2802-2817 (2009).
86. Abdul-Majeed, S., Moloney, B.C. & Nauli, S.M. Mechanisms regulating cilia growth and cilia function in endothelial cells. *Cellular and molecular life sciences : CMLS* **69**, 165-173 (2012).

9 LIST OF PUBLICATIONS

- Compartmentalized cAMP Signaling in Neurodegenerative Diseases
M. Sepe, L. Rinaldi, M. Porpora, R. Delle Donne, **S. Sauchella** And A. Feliciello. European Journal Of Neurodegenerative Diseases. Vol. 3, No. 3, 119-128 (2014)
- **Sauchella S***, Porpora M*, Rinaldi L., Sepe M., Delle Donne R., Torres-Quesada O., Intartaglia D., Garbi C., Insabato L., Santoriello M., Bachmann V.A., Synofzik M., Ivan C., Stefan E., Feliciello A. (2017) Counterregulation of cAMP-directed kinase activities controls ciliogenesis. *Submitted for publication. Currently, under minor revision.*
***Equally contributed.**