BIBLIOGRAFIA

AA.VV. (1985). Seminario informativo sui risultati del Progetto Laghi Costieri. Provincia di Latina. Latina.

AGAWIN S.R., DUARTE C.M., AGUSTÍ S. (2000). Nutrient and temperature control of the contribution of picoplankton to phytoplankton biomass and production. *Limnol. Oceanogr.* 45(3): 591-600.

ANDERSSON A., HAECKY P., HAGTRÖM Å. (1994). Effect of temperature and light on the growth of micro- nano- and pico-plankton: impact on algal succession. *Marine Biology* 120: 511-520.

BASSET A. AND ABBIATI M. (2004). Challenges to transitional water monitoring:ecological descriptor and scales. *Aquatic. Conserv.: Mar. Freshw. Ecosyst.* 14: S1-S3.

BASSET A., SANGIORGIO F., PINNA M. (2004). Monitoring with benthic macroinvertebrate: advantage and disadvantages of body size descriptors. *Aquatic. Conserv.: Mar. Freshw. Ecosyst.* 14: S43-S58.

BASSET A., SABETTA L., CARRADA G.C. (2006). Conservation of transitional water ecosystems in Mediterranean area: bridging basic ecological research and theories with requirements of application. *Aquatic Conserv: Mar. Freshw. Ecosyst.*16: 439-440.

BASSET A., SABETTA L., FONNESU A., MOUILLOT D. , DO CHI T., VIAROLI P. , REIZOPOULOU S., CARRADA G.C. (2006). Typology in Mediterranean Transitional Waters: new challenges and perspectives. *Aquatic Conserv: Mar. Freshw. Ecosyst.*16: 441-455.

BEC B., HUSSEINI-RATREMA J., COLLOS Y., SOUCHU P., VAQUER A. (2005). Phytoplankton seasonal dynamics in a Mediterranean coastal lagoon: emphasis on the picoeukaryote community. *J. Plankton Research* 27(9): 881-894.

BELL T. AND KALFF J. (2001). The contribution of picophytoplankton in marine and freshwater systems of different trophic status and depth. *Limnol. Oceanogr.* 46(5): 1243-1248.

BIANCHI C.N. (1988). Tipologia ecologica delle lagune costiere italiane. In: *Le lagune costiere: ricerca e gestione,* Carrada G.C., Cicogna F. e Fresi E. (eds.), CLEM Pubbl.: 57-66.

BINDER B.J., CHISHOLM S. W:, OLSON R. J. FRANKEL S.L. AND WORDEN A.Z. (1996). Dynamics of picophytoplankton, ultraplankton and bac teria in the central equatorial Pacific. *Deep Sea Research II* 43:907-931

BLOMQVIST P. PETTERSSON A. AND HYENSTRAND P. (1994). Ammonium-nitrogen: a key regulatory factor causing dominance of non-nitrogen-fixing cyanobacteria in aquatic systems. *Arch. Hydrobiologie* 132:141-164

BORJA A., FRANCO J., PEREZ V. (2000). A marine Biotic Index to establish the ecological quality of soft-bottom benthos within European estuarine and coastal environments . *Mar. Poll. Bull.* 40:1100-1114.

BRAMBATI A. (1988). Lagune e stagni costieri: due ambienti a confronto. In: *Le lagune costiere: ricerca e gestione,* Carrada G. C., Cicogna F. e Fresi E. (eds.), CLEM Pubbl.: 9-33.

CARRADA G.C. (2004). Editorial. *Uomo e Natura* 2: 3.

CARRADA G.C., GRIECO M., TALAMO A., MANGONI O. Presence and role of picophytoplankton in transitional waters. *Tansitional Water Bullettin* (submitted).

CAROPPO C. (2000). The contribution of picophytoplankton to community structure in a Mediterranean brackish environment. *J. Plankton Research* 22(2): 381-397.

CHAPELLE A., MENESGUEN A., DESSOUS-PAOLI J.-M., SOUCHU P., MAZOUNI N., VAQUER A., MILLET B. (2000). Modelling nitrogen, primary production and oxygen in a Mediterranean lagoon. Impact of oysters farming and inputs from the watershed. *Ecological Modelling* 127(2-3): 161-181.

CHISHOLM S.W. (1992). Phytoplankton size. In *Primary productivity and biogeochemical cycles in the sea*, Falkowski PG, Woodhead AD (Eds). Plenum Press, New York.

COLLOS Y., HUSSEINI-RATREMA J., BEC B., VAQUER A., LAM HOAI T., ROUGIER C., PONS V., SOUCHU P. (2005). Pheopigment dynamics, zooplankton grazing rates and the autumnal ammonium peak in a Mediterranean lagoon. *Hydrobiologia* 550(1):83-93.

CONNELL J.H.(1978). Diversity in tropical rain forests and coral reefs. *Science* 199:1302-1310.

CHRISTIAN R.R. (2005). Beyond the Mediterranean to global observations of coastal lagoons. *Hydrobiologia* 550: 1-8.

DE CASABIANCA M.-L., LAUGIER T., MARINO-SORIANO E. (1997). Seasonal changes of nutrients in water and sediment in a Mediterranean lagoon with shellfish farming activity (Thau lagoon, France). *ICES Journal of Marine Science* 54: 905-916.

DE PIPPO T., DANADIO C., PENNETTA M. (2002-2002). Evoluzione morfologica della Laguna di Sabaudia (Mar Tirreno, Italia Centrale). *Geologia Romana*, 36: 1-12.

DE ROOS A.M., PERSONN L., MCCAULEY E. (2003). the influence of size-dependent life-history traits on the structure and dynamics of populations and communities. *Ecology Letters* 6: 473-487.

FANO E.A., MISTRI M., ROSSI R. (2003). The ecofunctional quality index (EQI): a new tool for assessing lagoonal ecosystem impairment. *Estuarine Coastal and Shelf Science*, 56: 709-716.

FOUILLAND E., DESCOLAS-GROS C., COLLOS Y., VAQUER A., SOUCHU P., GASC A., BIBENT B., PONS V. (2002). Influence of nitrogen enrichment on size-fractionated in vitro carboxylase activities of phytoplankton from Thau Lagoon (Coastal Mediterranean Lagoon, France). *Journal of Experimental Marine Biology and Ecology* 275: 147-171.

FERRERO L. (1961). Ricerche chimico-fisiche e biologiche sui laghi salmastri pontini in relazione alla produttività. Il lago di Paola (Sabaudia). Ricerche quantitative sulla fauna bentonica. *Boll. Pesca Pescicoltura Idrobiologia* 16: 173-203.

FRONEMAN P.W., PERNISSINOTTO R., MCQUAID C.D. (1996). Seasonal variation in microzooplankton grazing in the region of the subtropical Convergence. *Marine Biology*, 126: 433-442.

GIESKES W.W. AND KRAAY G.W. (1986) Floristic and physiological differences between shallow and the deep nanophytoplankton communities in the euphotic zone of the trpical Atlantic ocean revealed. By HPLC analysis of pigments. *Nature* 91: 567-576

GILABERT J. (2001). Seasonal plankton dynamics in a Mediterranean hypersaline coastal lagoon: the Mar Menor. *J. Plankton Research* 23(2): 207-217.

GILABERT J. (2001). Short-term variability of the planktonic size structure in a Mediterranean coastal lagoon. *J. Plankton Research* 23(2): 219-226.

GILBERT F., SOUCHU P., BIANCHI M., BONIN P. (1997). Influence of shellfish farming activities on nitrification, nitrate reduction to ammonium and denitrification at the water-sediment interface of Thau lagoon, France. *Mar. Ecol. Prog. Ser.* 151: 143-153.

GUELORGET O. ET PERTHUISOT J.P., (1983). Le domain paralique. *Trav. Lab. Géol.*, 16:1-136, École Norm. Sup., Paris.

HEIN M., PEDERSEN M.F. AND SAND.JENSEN K. (1995) Size-dependent nitrogen uptake in micro- and macroalgae. *Marine Ecology Progr. Ser.* 118: 247-253

HELBLING E.W., VILLAFAÑE V., HOLM-HANSEN O.(1991). Effect of iron on productivity and size distribution of Antarctic phytoplankton. *Limnol. Oceanogr.* 36:1879-1885.

HLAILI SAKKA A., GRAMI B., MABROUK HADJ H. (2007). Phytoplankton growth and microzooplankton grazing rates in a restricted Mediterranean lagoon (Bizerte Lagoon, Tunisia). *Mar. Biol.* 151: 767-783.

HLAILI SAKKA A., GRAMI B., NIQUIL N., GOSSELIN M., HAMEL D., TROUSSELLIER M., MABROUK HADJ H. (2008). The planktonic food web of Bizerte lagoon (south-western Mediterranean) during summer: I. Spatial distribution under different anthropogenic pressures. *Estuarine Coastal and Shelf Science* 78(1): 61-77.

HOLLING C.S. (1992). Cross-scale morphology, geometry and dynamics of ecosystems. *Ecological Monographs*, 62: 447-502

HOLM-HANSEN O., LORENZ C.J., HOLMES R.W., STRICKLAND J.D.H.(1965). Fluorimetric determination of chlorophyll. *J. Cons. Perm. Int. Explor.* Mer., 30:3-15.

INNAMORATI M., FERRARI I., MARINO D., RIBERA D'ALCALÀ M. (Ed.s) (1990). Metodi nell'ecologia del plancton marino. *Nova Thalassia* 11: 372 pp

INTERNATIONAL UNION OF BIOLOGICAL SCIENCES (1959). Simposium on the classification of brackish waters – series B, N°35. Paris.

IRIARTE J.L., GONZALEZ H.E. (2004). Phytoplankton size structure during and after the 1997/98 El Niño in a coastal upwelling area of the northern Humboldt Current System. *Mar. Ecol. Pro. Ser.* 269:83-90.

JEFFREY S. W. AND MANTOURA R. F. C. (1997). Development of pigment methods for oceanography: SCOR-supported Working Groups and objectives. In: *Phytoplankton pigments in oceanography: guidelines to modern methods,* Jeffrey S.W., Mantoura R.F.C., Wrigth S.W. (Eds) UNESCO. Publishing, pp. 19-36.

JOCHEM F. (1988). On the distribution and importance of picocyanobacteria in a boreal inshore area (Kiel Bight, western Baltic). *J. Plankton Research* 10:1009-1022

KNOWLTON N. (1993). Sibling species in the sea. *Ann. Rev. Ecol. Syst.* 24:189-216.

KUOSA H. (1991). Picoplankton algae in the northern Baltic Sea: seasonal dynamics and flagellates grazing. *Mar. Ecol. Progr. Ser.*, 73: 269-276.

LA MURA V. (1995). Caratterizzazione idrodinamica della laguna di Sabaudia (Mar Tirreno) ed implicazioni sulle metodologie di campionamento del particellato. Tesi di dottorato, VIII ciclo, Università di Messina.

LARDICCI C. AND ROSSI F. (1998). Detection of stress on macrozoobenthos: evaluation of some methods in a coastal Mediterranean lagoon. *Mar. Environ. Res.* 45:367-386.

LAWRENCE D., DAGG M.J., LIU H.B., CUMMINGS S.R., ORTNER P.B., KELBLE C. (2004). Wind events and benthic-pelagic coupling in a shallow subtropical bay in Florida. *Mar. Ecol. Progr. Ser.* 266:1-13.

LE GALL S., BEL HASSEN M., LE GALL P. (1997). Ingestion of a bacterivorous ciliate by the oyster *Crassostrea gigas*: protozoa as a trophic link between picoplankton and benthic suspension-feeders. *Mar. Ecol. Prog. Ser.* 152: 301-306.

LEVIN L.A., BOESCH D.F., COVICH A., DAHM C., ERSEUS C., EWEL K.C., KNEIB R.T., MOLDENKE A., PALMER M.A., SNELGROVE P., STRAYER D., WESLAWSKI J.M. (2001). The function of marine critical transition zones and importance of sediment biodiversity. *Ecosystems* 4:430-451.

LI W.K.W. (1998). Annual average abundance of heterotrophic bacteria and Synecococcus in surface ocean waters. *Limnol. Oceanogr*, 43:1746-1753.

LÓPEZ-FLORES R., BOIX D., BADOSA A., BRUCET S., QUINTANA X.D. (2006). Pigment composition and size distribution of phytoplankton in a confined Mediterranean salt marsh ecosystem. *Marine Biology* 149: 1313-1324.

MAGAZZÙ G., DECEMBRINI F. (1995). Primary production, biomass and abundance of phototrophic picoplankton in the Mediterranean Sea: a rew. *Aquatic. Microb. Ecol.* 9: 97:104.

MANTOURA R.F.C. AND LLEWELLYN C.A. (1983). The rapid determination of algal chlorophyll and carotenoid pigments and their breakdown products in natural waters by reverse phase HPLC. *Anal. Chim. Acta*, 151: 297- 314.

MINE BERG G., BALODE M., PURINA I., BECERE S., BÉCHEMIN C., MAESTRINI S.Y. (2003). Plankton community composition in relation to availability and uptake of oxidized and reduced nitrogen. *Aquat. Microb. Ecol.* 30: 263-274.

MIRACLE M.R., GARCÍA M.P., VICENTE E. (1984). Heterogeneidad espacial de las comunidades fitoplanctonicas de la Albufera de Valencia. *Limnética* 1:20-31.

MOON-VAN DER STAAY S.Y., MOON-VAN DER STAAY G.W.M., GUILLOU L., VAULOT D., CLAUSTRE H., MEDLIN L.K. (2000). Abundance and diversity of prymnesiophytes in the picoplankton community from equatorial Pacific Ocean inferred from 18S rDNA sequences. *Limnol. Oceanogr.* 45(1): 98-109.

MOORE L.R., POST A.F. ROCAP G., CHISHOLM S.W. (2002). Utilization of different nitrogen sources by the marine cyanobacteria *Prochlorococcus* and *Synechococcus*. *Limnol. Oceanogr.*, 47:989-996.

MOREIRA-TURCQ P.F., CAUWET G., MARTIN J.M. (2001). Contribution of flow cytometry to estimate picoplankton biomass in estuarine systems. *Hydrobiologia* 462: 157-168.

MOUILLOT D., CULIOLI J.M., DO-CHI T., (2002). Indicator species analysis as a test of non-random distribution of species in the context of marine protected areas. *Environ. Conserv.* 29:385-390.

MOUILLOT D., SPATHARIS S., REIZOPOULOU S., LAUGIER T., SABETTA L., BASSET A., DO-CHI T., (2006). Alternatives to taxonomic-based approaches to assess changes in transitional water communities. *Aquat. Conserv.:Mar. Freshw. Ecosyst.* 16: 469-482.

MURRELL M.C. AND HOLLIBAUGH J.T. (1998). Microzooplankton grazing in northern San Francisco Bay measured by the dilution method. *Aquatic Microbial Ecology* 15: 53-63.

NICHOLS S., WEBER S., SHAW B., (2000). A proposed aquatic plant community biotic index for Wisconsin lakes. *Environ. Manag,* 26: 491-502.

OVIATT C., LANE P., FRENCH F.III, DONAGHAY P. (1989). Phytoplankton speciesand abundance in response to eutrophication in coastal marine mesocosms. *J. Plankton Research*, 11: 1223-1244.

PAINTING S.J., MOLONEY C.L., LUCAS M.I. (1993). Simulation and field measurements of phytoplankton-bacteria-zooplankton interactions in the southern Benguela upwelling region. *Mar. Ecol. Progr. Ser.* 100:55-69.

PARTENSKY F., BLANCHOT J., LANTOINE F., NEVEUX J., MARIE D. (1996). Vertical structure of picoplankton at different trophic sites of tropical northeastern Atlantic Ocean. *Deep-Sea Res.*I,43(8): 1191-1213.

PERDICARO R. (1983). Osservazioni sulle condizioni ideologiche del lago di Sabaudia, o di Paola nel biennio 1982-83. *Quaderni Ist. Idrobiologia ed Acquacoltura Brunelli* 3(2): 59-72.

PERDICARO R. (1984). Condizioni chimico-fisiche del lago di Sabaudia o di Paola dal 1933 al 1984. *Quaderni Ist. Idrobiologia ed Acquacoltura Brunelli* 4(2): 71-139.

QUINONES R.A. (1994). A comment on the use of allometry in the study of pelagic ecosystem. In *The size structure and metabolism of the pelagic ecosystem*. Rodríguez J., Li WKW (eds). *Scientia marina*, 58: 11-16.

QUINTANA X.D., MORENO-AMICH R., COMÍN F.A. (1998). Nutrient and plankton dynamics in a Mediterranean salt marsh dominated by incidents of flooding. Part 1: differential confinement of nutrients. *J. Plankton Research* 20(11): 2089-2107.

QUINTANA X.D. AND MORENO-AMICH R. (2002). Phytoplankton composition of Empordà salt marshes, Spain and its response to freshwater flux regulation. *J. Coastal Research* SI 36: 581-590.

QUINTANA X.D., COMÍN F.A., MORENO-AMICH R. (2002). Biomass-size spectra in aquatic communities in shallow fluctuating Mediterranean salt marshes (Empordà wetlands, NE Spain). *J. Plankton Research* 24(11): 1149-1161.

RAVEN J.A. (1998). The twelfth Tansley lecture. Small is beautiful: the picophytoplankton. *Functional Ecology* 12: 503-513.

REIZOPOULOU S., THESSALOU-LAGAKI M., NICOLAIDOU A. (1996). Assessment of disturbance in Mediterranean lagoons: an evaluation of methods. *Mar. Biol.* 125:189-197.

REIZOPOULOU S. AND NICOLAIDOU A. (2005). Index of Size Distribution (ISD): a method for assessing disturbance in lagoons. In *International Conference on Lagoons and Coastal Wetlands in Global Change Context: Impacts and Management Issues*, UNESCO Proceeding Series, Venice, in press.

ROBSON B.J., BARMUTA L.A., FAIRWEATHER P.G. (2005). Methodological and conceptual issues in the search for a relationship between animal body-size distribution and benthic habitat architecture. *Marine and Freshwater Research* 56: 1-11.

RODRÌGUEZ J., JIMÉNEZ F., BAUTISTA B., RODRÌGUEZ V. (1987). Planktonic biomass spectra dynamics during a winter production pulse in Mediterranean coastal waters. *J. Plankton Research* 9: 1183-1194.

RODRÌGUEZ J., BLANCO J.M., JIMÉNEZ-GOMEZ F., ECHEVARRÍA F., GIL J., RODRÌGUEZ V., RUIZ J., BAUTISTA B., GUERRERO F. (1998). Patterns in the size structure of the phytoplankton community in the deep fluorescence maximum of the Alborean Sea (Southwestern Mediterranean). *Deep-Sea Res.* 45: 1577-1593.

RODRÌGUEZ J., TINTORÉ J., ALLEN J.T., BLANCO J.M., GOMIS D., REUL A., RUIZ J., RODRÌGUEZ V., ECHEVARRÍA F., JIMÉNEZ-GOMEZ F., (2001). Mesoscale vertical motion and size structure of phytoplankton in the ocean. *Nature* 410: 360-363.

RODRÌGUEZ J., JIMÉNEZ-GOMEZ F., BLANCO J.M., FIGUEROA F.L. (2002). Physical gradients and spatial variability of the size structure and composition of phytoplankton in the Gerlache Strait (Antarctica). *Deep-Sea Res.* II 49: 693-706.

SCHMIDT D.N., THIERSTEIN H. R., BOLLMANN J., SCHIEBEL R. (2004). Abiotic forcing of plankton evolution in the cenozoic. *Science*, 303: 207-210.

SELMER J.S., FERRIER-PAGES C., CELLARIO C., RASSOULZADEGAN F. (1993). New and regenerated production in relation to the microbial loop in the NW Mediterranean Sea. *Mar. Ecol. Prog. Ser.* 100:71-83.

SERRA T., GRANATA T., COLOMER J., STIPS A., MØHLENBERG F., CASAMITJANA X. (2003). The role of advection and turbulent mixing in the vertical distribution of phytoplankton . *Estuar. Coast. Shelf. Sci.* 56: 53-62.

SIN Y., WETZEL R.L., ANDERSON I.C. (2000). Seasonal variation of size-fractionated phytoplankton along the gradient in the York River estuary, Virginia (USA). *J. Plankton Research* 22(10): 1945-1960.

SOROKIN P.YU., SOROKIN YU.I., BOSCOLO R., GIOVANARDI O. (2004). Bloom of picocyanobacteria in the Venice lagoon during summer-autumn 2001: ecological consequences. *Hydrobiologia* 523: 71-85.

SOUCHU P., VAQUER A., COLLOS Y., LANDREIN S., DESLOUS-PAOLI J.M., BIBENT B. (2001). Influence of shellfish farming activities on the biogeochemical composition of water column in thau lagoon. *Mar. Ecol. Progr. Ser.* 218: 141-152.

TAKAHASHI M. AND BIENFANG, P.K. (1983). Size structure of phytoplankton biomass and photosynthesis in subtropical Hawaiian waters. *Mar. Biol.*, 76: 203-211.

VAQUER A., TROUSSELLIER M., COURTIES C., BIBENT B. (1996). Standing stock and dynamics of picophytoplankton in the Thau Lagoon (northwest Mediterranean coast). *Limnol. Oceanogr.* 4(8):1821-1828.

VELDHUIS M.J.W., KRAAY G.W., VAN BLEIJSWIJK J.D.L., BAARS M.A. (1997). Seasonal and spatial variability in phytoplankton biomass, productivity and grow in the north-western Indian Ocean: the southwest and northeast monsoon, 1992-1993. *Deep-Sea Res. I* 44: 425-449.

VELDHUIS M.J.W., TIMMERMANS K.R., CROOT P., VAN DER WAGT B. (2005). Picophytoplankton; a comparative study of their biochemical composition and photosynthetic properties. *J. Sea Research* 53: 7-24.

VIDUSSI F., CLAUSTRE H., BUSTILLOS-GUZMAN J., CAILLIAU C., MARTY J.-C. (1996). Determination of chlorophylls and carotenoids of marine phytoplankton: separation of chlorophyll a from divinyl-chlorophyll a and zeaxanthin from lutein. *J. Plankton Research* 18: 2377-2382.

WALSH J.J. (1976). Herbivory as a factor in patterns of nutrient utilization in the sea. *Limnol. Oceanogr.*, 21: 1-13.

WFD 2000. European Commission Directive 2000/60/EC of European Parliament and of Council of 23 October 2000, establishing a framework for Community action in the field of water policy. *Official Journal of the European Communities*, L 327/1-72, Brussels.